


The Law of Sines


Solving a triangle means finding the measures of all the sides and angles. An **oblique triangle** is a triangle without a right angle. To solve an oblique triangle, we must know three pieces of information, at least one of which must be the length of a side. (Three angles define an infinite number of triangles).



The Law of Sines: In any triangle, $\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}$


ASA or AAS: Find the third angle using the fact that the three angles of a triangle add to 180° . Then use the law of sines to find the other two sides of the triangle.

Example: $\alpha = 28^\circ$, $\beta = 66^\circ$, $c = 8.2$




SSA (The Ambiguous Case): If you know two sides and a non-included angle (an angle that is not between the sides), there may be zero, one, or two possible triangles that fit the given measurements. To figure out how many triangles there are for an *acute* angle α , do the following:


1. Draw the given angle (α) in standard position with a terminal side of length b . Don't draw side a yet.
2. Let h be an altitude from C to the initial side of α .




3. Since $\sin \alpha = h/b$, $h = b \sin \alpha$. Compare h to a as follows:
 - a. If $a < h$, then no triangle can be formed.
 - b. If $a = h$, then one triangle (a right triangle) can be formed.
 - c. If $h < a < b$, then two triangles can be formed.
 - d. If $a \geq b$, then one triangle can be formed.





(a) $a < h$
no triangle



(b) $a = h$
one triangle





(c) $h < a < b$
two triangles



(d) $a \geq b$
one triangle

If α is *obtuse*, there is either no triangle (if $a \leq b$) or one triangle (if $a > b$) possible.



Examples:

a) $\beta = 38^\circ$, $b = 2.9$, $c = 5.9$

b) $\beta = 38^\circ$, $b = 6.4$, $c = 5.9$

c) $\beta = 38^\circ$, $b = 4.7$, $c = 5.9$

d) $\gamma = 125^\circ$, $b = 5.7$, $c = 8.6$


Bearing: The measure of an angle that describes the direction of a ray is called the bearing. Bearing is the clockwise angle from due north.

Another way to express bearing is to describe the acute angle that the ray makes with a ray pointing due north or south. For example:


N 60° E is a bearing of 60° east of north

S 30° E is a bearing of 30° east of south

S 45° W is a bearing of 45° west of south



Example: During an important NATO exercise, an F-14 Tomcat left the carrier Nimitz on a course with a bearing of 34° and flew 400 miles. Then the F-14 flew for some distance on a course with a bearing of 162° . Finally, the plane flew back to its starting point on a course with a bearing of 308° . What distance did the plane fly on the final leg of the journey? Round to the nearest tenth of a mile.

The Law of Cosines

$$a^2 = b^2 + c^2 - 2bc \cos \alpha$$

The Law of Cosines: In any triangle, $b^2 = a^2 + c^2 - 2ac \cos \beta$

$$c^2 = a^2 + b^2 - 2ab \cos \gamma$$

SSS: Use the fact that the largest angle is across from the longest side of the triangle to solve for the largest angle using the law of cosines. (For example, if c is the longest side, use the equation $c^2 = a^2 + b^2 - 2ab \cos \gamma$ to solve for γ .) Then use the law of sines to find the remaining angles, which will both be acute.

Example: $a = 3.8$, $b = 9.6$, $c = 7.7$


SAS: Find the length of the third side using the law of cosines. Use the law of sines to find the angle across from the shorter of the two given sides. Find the remaining angle by subtracting the first two from 180° .

Example: $b = 5.8$, $c = 3.6$, $\alpha = 39.5^\circ$

Example: Jan and Dean started hiking from the same location at the same time. Jan hiked at 4 mph with bearing N12°E, and Dean hiked at 5 mph with bearing N31°W. How far apart were they after 6 hours? Round to the nearest tenth of a mile.

Length of a Chord: If a chord of length a is intercepted by a central angle α in a circle of radius r , then $a = r\sqrt{2 - 2\cos\alpha}$. (This formula is derived from the law of cosines.)

Example: Find the length of the chord intercepted by a central angle of 33.8° in a circle of radius 22.4 ft.



Area of a Triangle


The formula $A = \frac{1}{2}bh$ gives the area of a triangle.

In the diagram at the right, $\sin \alpha = \frac{h}{c}$, so $h = c \sin \alpha$. Using

substitution, we derive the formula $A = \frac{1}{2}bc \sin \alpha$.

Depending on which angles and sides are known, the

formulas $A = \frac{1}{2}ac \sin \beta$ and $A = \frac{1}{2}ab \sin \gamma$ can also be used.



Using the law of cosines, it is possible to derive a formula for the area of a triangle that involves only the lengths of the sides of the triangle. The formula is known as “Heron’s Formula” after Heron of Alexandria, who is believed to have discovered it around AD 75.

Heron’s Formula: The area of a triangle with sides of lengths a , b , and c is given by:

$$A = \sqrt{S(S-a)(S-b)(S-c)}, \text{ where } S = (a+b+c)/2.$$

Examples:

Find the area of the triangle with $\alpha = 39.4^\circ$, $b = 12.6$, and $c = 13.7$

Find the area of a triangle with $\alpha = 56.3^\circ$, $\beta = 41.2^\circ$, and $a = 9.8$

Find the area of the triangle with $a = 12$, $b = 8$, and $c = 6$

Find the area of a triangle with $a = 346$, $b = 234$, and $c = 422$

Vectors

Scalar Quantities: Quantities such as length, area, volume, temperature, and time, which have magnitude (size), but no direction.

Vector Quantities: Quantities that involve both a magnitude and direction, such as velocity, acceleration, and force. These quantities can be represented by **directed line segments** called **vectors**.


The length of a vector represents the **magnitude** of the vector quantity. The **direction** is indicated by the position of the vector and the arrowhead at one end.

Notation: \overrightarrow{AB} is used to name a vector with **initial point** A and **terminal point** B . Vectors may also be denoted by bold letters. \overrightarrow{AB} can also be written as \mathbf{AB} . If the initial and terminal points are not specified, vectors can be named by a single uppercase or lowercase letter (eg. \vec{b} , \vec{B} , \mathbf{b} , or \mathbf{B} .) The magnitude of vector \mathbf{A} is written $|\mathbf{A}|$.


Equal Vectors: Vectors with the same magnitude and direction. They do not have to be in the same place.

Zero Vector: A vector with no magnitude and no direction. It is denoted by $\mathbf{0}$.

Scalar Multiplication: For any scalar k and vector \mathbf{A} , $k\mathbf{A}$ is a vector with magnitude $|k|$ times the magnitude of \mathbf{A} . If $k > 0$, then the direction of $k\mathbf{A}$ is the same as the direction of \mathbf{A} . If $k < 0$, the direction of $k\mathbf{A}$ is opposite to the direction of \mathbf{A} . If $k = 0$, then $k\mathbf{A} = \mathbf{0}$.




Suppose two draft horses are pulling on a tree stump with forces of 200 pounds and 300 pounds, with an angle of 65° between the forces. If \mathbf{A} and \mathbf{B} had the same direction, then there would be a total force of 500 pounds acting on the stump, but the total force is less because of the angle between the forces. By the **parallelogram law**, the force acting along the diagonal of the parallelogram, with a magnitude equal to the length of the diagonal, has the same effect on the stump as the two forces \mathbf{A} and \mathbf{B} . The force $\mathbf{A} + \mathbf{B}$ acting along the diagonal is called the **sum** or **resultant** of \mathbf{A} and \mathbf{B} .



Vector Addition: To find the resultant or sum $\mathbf{A} + \mathbf{B}$ of any vectors \mathbf{A} and \mathbf{B} , position \mathbf{B} (without changing its magnitude or direction) so that the initial point of \mathbf{B} coincides with the terminal point of \mathbf{A} . The vector that begins at the initial point of \mathbf{A} and ends at the terminal point of \mathbf{B} is the vector $\mathbf{A} + \mathbf{B}$. For every vector \mathbf{A} , there is a vector $-\mathbf{A}$, with the same magnitude as \mathbf{A} , but the opposite direction. For any two vectors \mathbf{A} and \mathbf{B} , $\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$.


Example: Sketch the vectors $\mathbf{A} + \mathbf{B}$ and $\mathbf{A} - \mathbf{B}$.



Any nonzero vector \mathbf{w} is the sum of a **horizontal component**, w_x , and a **vertical component**, w_y . If a vector \mathbf{w} is placed in a rectangular coordinate system so that its initial point is the origin, then \mathbf{w} is called a **position vector**. The angle θ formed by the positive x -axis and a position vector is the **direction angle** for the position vector.

If the vector \mathbf{w} has magnitude r , direction angle θ , horizontal component w_x , and vertical component w_y , then we get

$$\cos \theta = \frac{|w_x|}{r} \text{ and } \sin \theta = \frac{|w_y|}{r} \text{ or } |w_x| = |r \cos \theta| \text{ and } |w_y| = |r \sin \theta|.$$




Examples: Find the magnitude of the horizontal and vertical components for each vector \mathbf{v} with the given magnitude and direction angle θ . Round to the nearest tenth.

a) $|\mathbf{v}| = 5.6$, $\theta = 22^\circ$

b) $|\mathbf{v}| = 445$, $\theta = 211.1^\circ$

Component Form: The notation $\langle x, y \rangle$ is used to define a position vector with terminal point (x, y) . This is called component form because the horizontal component is $\langle x, 0 \rangle$ and its vertical component is $\langle 0, y \rangle$.

The magnitude of the vector $\mathbf{v} = \langle x, y \rangle$ is $|\mathbf{v}| = r = \sqrt{x^2 + y^2}$. To find the direction angle, use $\sin \theta = \frac{y}{r}$, $\cos \theta = \frac{x}{r}$, and $\tan \theta = \frac{y}{x}$.



If a vector has magnitude r and direction angle θ , its component form is $\langle r \cos \theta, r \sin \theta \rangle$.

Examples: Find the magnitude and direction angle of each vector.

a) $\mathbf{v} = \langle 2, -6 \rangle$

b) $\mathbf{v} = \langle -3, 2 \rangle$

c) $\mathbf{v} = \langle -4, -5 \rangle$

Examples: Find the component form for each vector \mathbf{v} with the given magnitude and direction angle θ .

a) $|\mathbf{v}| = 12$, $\theta = 45^\circ$

b) $|\mathbf{v}| = 50$, $\theta = 120^\circ$

If $\mathbf{A} = \langle a_1, a_2 \rangle$, $\mathbf{B} = \langle b_1, b_2 \rangle$, and k is a scalar, then

	1. $k\mathbf{A} = \langle ka_1, ka_2 \rangle$	Scalar Product
Vector Arithmetic:	2. $\mathbf{A} + \mathbf{B} = \langle a_1 + b_1, a_2 + b_2 \rangle$	Vector Sum
	3. $\mathbf{A} - \mathbf{B} = \langle a_1 - b_1, a_2 - b_2 \rangle$	Vector Difference
	4. $\mathbf{A} \cdot \mathbf{B} = a_1b_1 + a_2b_2$	Dot Product

Examples: Let $\mathbf{w} = \langle -1, -3 \rangle$ and $\mathbf{v} = \langle -3, 4 \rangle$. Perform the operations indicated.

a) $\mathbf{w} - \mathbf{v}$

b) $-8\mathbf{v}$

c) $3\mathbf{w} + 4\mathbf{v}$

d) $\mathbf{w} \cdot \mathbf{v}$

The Angle Between Two Vectors:

If \mathbf{A} and \mathbf{B} are nonzero vectors and α is the smallest positive angle between them, then $\cos \alpha = \frac{\mathbf{A} \cdot \mathbf{B}}{|\mathbf{A}||\mathbf{B}|}$.

Examples: Find the smallest positive angle between the following vectors:

a) $\langle 1, 3 \rangle$ and $\langle 5, 2 \rangle$

b) $\langle -5, 1 \rangle$ and $\langle 7, 3 \rangle$

Parallel Vectors: The vectors \mathbf{A} and \mathbf{B} are parallel if and only if $\mathbf{A} = k\mathbf{B}$ for a nonzero scalar k .

Perpendicular Vectors: The vectors \mathbf{A} and \mathbf{B} are perpendicular if and only if $\mathbf{A} \cdot \mathbf{B} = 0$.

Examples: Determine whether each pair of vectors is parallel, perpendicular, or neither.

a) $\langle -2, 3 \rangle$ and $\langle 6, 4 \rangle$

b) $\langle 2, -5 \rangle$ and $\langle -4, 10 \rangle$

c) $\langle 2, 6 \rangle$ and $\langle 6, 2 \rangle$

The vectors $\mathbf{i} = \langle 1, 0 \rangle$ and $\mathbf{j} = \langle 0, 1 \rangle$ are called **unit vectors** because each has magnitude one. For any vector $\langle a_1, a_2 \rangle$, we have $\langle a_1, a_2 \rangle = a_1 \langle 1, 0 \rangle + a_2 \langle 0, 1 \rangle = a_1\mathbf{i} + a_2\mathbf{j}$. The form $a_1\mathbf{i} + a_2\mathbf{j}$ is called a **linear combination** of the vectors \mathbf{i} and \mathbf{j} .

Examples: Write each vector as a linear combination of the unit vectors \mathbf{i} and \mathbf{j} .

a) $\mathbf{A} = \langle 2, 3 \rangle$

b) $\mathbf{B} = \langle -1, 7 \rangle$


c) $\mathbf{C} = \langle 0, -9 \rangle$

Applications of Vectors

Example: Forces of 2 lb and 12 lb are acting at an angle of 72° to each other. Find the magnitude of the resultant force and the angle between the resultant and each force.

Example: The resultant of a 13-lb force and another force has a magnitude of 9 lb at an angle of 20° with the 13-lb force. Find the magnitude of the other force and the angle between the two forces.

Inclined Plane Problems: The weight of an object is always modeled as a vertical vector and the force required to move the object is modeled as a vector parallel to the inclined plane. Its length increases as the incline increases. The resultant of these two forces is a vector perpendicular to the inclined plane. It is what a bathroom scale would read if trapped between the object and the plane.




Example: Find the amount of force required to push an 800-pound block of ice up a ramp that is inclined 10° .

Example: A landscaper uses 100 pounds of force to pull a cart full of rocks up a driveway that is inclined 15° . What is the weight of the cart?

Example: If 300 pounds of force is required to push a 1000-pound safe up a ramp, then what is the angle of inclination of the ramp?

Navigation Problems: Wind affects the speed and direction of a plane. The **heading** and **air speed** are the direction and speed of the plane before wind is taken into account. The **course** and **ground speed** are the direction and speed of the plane with wind taken into account. The angle between the heading and the course is the **drift angle**.



Example: An airplane is headed due east with an air speed of 200 mph. The wind is out of the south (bearing 0°) at 40 mph. Find the ground speed, the drift angle, and the course of the airplane.

Example: An airplane is headed due west with an air speed of 400 mph. The wind is out of the southwest (bearing 45°) at 90 mph. Find the ground speed, the drift angle, and the course of the airplane.

Example: A boat is traveling at 20 mph with a bearing of $N20^\circ W$. The current is moving at 4mph with a bearing of $S40^\circ W$. Find the boat's true speed, drift angle, and course.